



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hence ten days on chronometer C equals 10 days, 1 hour, 12 minutes and 55 seconds on chronometer A . Therefore the time by the true watch A is 12 minutes, 55 seconds past 1 P. M. on the tenth day.

Also solved by CHARLES C. CROSS, and G. B. M. ZERR.

PROBLEMS FOR SOLUTION.

ALGEBRA.

90. Proposed by J. MARCUS BOORMAN, Consultative Mechanician and Counselor at Law, Woodmere, Long Island, N. Y.

Fully solve $x^2 + y^2 + m(x+y) = m^2 \dots\dots (I)$; $x^2 + y^2 + xy = m^2 \dots\dots (II)$; give process, roots, and corollary. [Solved in part, *Cirotte's Algebra*, page 202.]

91. Proposed by NELSON S. RORAY, Professor of Mathematics, South Jersey Institute, Bridgeton, N. J.

Solve the following without making use of the determinant notation and prove that the results obtained are the roots.

$$10x - 2y + 4z = 5,$$

$$3x + 5y - 3z = 7,$$

$$x + 3y - 2z = 2.$$

92. Proposed by W. F. BRADBURY, A. M., Head Master Latin School, Cambridge, Mass.

Find the sum to n terms of $1 + 3^3 + 5^3 + \dots\dots$ [From *Charles Smith's Elementary Algebra*, page 403.]

*** Solutions of these problems should be sent to J. M. Colaw not later than December 10.

GEOMETRY.

105. Proposed by B. F. FINKEL; A. M., M.Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Sind A, B, C, D vier harmonische Punkte und beschreibt man über dem Durchmesser AC einen Kreis, von welchem S ein beliebiger Punkt ist, so wird derjenige Kreisbogen, welcher innerhalb des Winkels BSD liegt, entweder von A oder von C Strecke PB halbirt. [*Reye's Geometrie der Lage*, page 191.]

106. Proposed by C. HORNUNG, A. M., Professor of Mathematics, Heidelberg University, Tiffin, Ohio.

Upon the sides of any triangle ABC let the equilateral triangles ABD, BCE , and CAF be described, and let their exterior sides produced intersect, BE and AF in K , DB and FC in L , and DA and EC in M . Prove DK, EL, FM , parallel.

107. Proposed by T. W. PALMER, A. M., Professor of Mathematics, University of Alabama.

Construct a triangle, given base, vertical angle and radius of inscribed circle.

*** Solutions of these problems should be sent to B. F. Finkel not later than December 10.